

Centraliser Dimension of Partially Commutative Groups ^{*†}

Andrew J. Duncan Ilya V. Kazachkov[‡]

Vladimir N. Remeslennikov[‡]

February 1, 2008

Abstract

In paper [7] we investigated the centraliser dimension of groups. In the current paper we study properties of centraliser dimension for the class of free partially commutative groups and, as a corollary, we obtain an efficient algorithm for computation of centraliser dimension in these groups.

1 Introduction

Partially commutative groups have been extensively studied in several different guises. They are variously known as right-angled Artin groups; trace groups; graph groups or locally free groups. For an introduction to this class of groups and a survey of the literature see [8].

In this paper we study the lattices of centralisers in partially commutative groups. Centraliser lattices, of various groups, have been investigated by numerous authors: for example [10], [20], [17], [13], [15], [16], [19], [12], [11], [4], [5], [1], [3], [18], [14] and [7]. In particular, a detailed account of results in the field can be found in V. A. Antonov's book [2]. Following [15] and [14], here we focus attention on the height of the lattice of centralisers of a group: which we call the centraliser dimension. Our interest in centralisers stems from their importance in constructing algebraic geometry over groups. We plan to apply the results of this paper to the study of, in the first instance, solutions of equations in one variable in partially commutative groups.

Partially commutative groups have been extensively studied in several different guises. They are variously known as right-angled Artin groups; trace groups; graph groups or locally free groups. For an introduction to this class of groups and a survey of the literature see [8].

^{*}Research supported by EPSRC grant GR/R87406/01

[†]2000 *Mathematics Subject Classification* 20F36 (primary), 20E15, 20F10 (secondary)

[‡]Supported by RFFI grant N05-01-00057-a

Let Γ be a finite, undirected, simple graph. Let $X = V(\Gamma)$ be the set of vertices of Γ and let $F(X)$ be the free group on X . For elements g, h of a group we denote the commutator $g^{-1}h^{-1}gh$ of g and h by $[g, h]$. Let

$$R = \{[x_i, x_j] \in F(X) \mid x_i, x_j \in X \text{ and there is an edge from } x_i \text{ to } x_j \text{ in } \Gamma\}.$$

We define the *free partially commutative group with (commutation) graph* Γ to be the group $G(\Gamma)$ with presentation $\langle X \mid R \rangle$.

The elements of X are termed the *canonical* generators of $G(\Gamma)$. In this article we refer to finitely generated free partially commutative groups as *partially commutative* groups.

If S is a subset of a group G then the centraliser of S in G is $C_G(S) = \{g \in G : gs = sg, \text{ for all } s \in S\}$. We write $C(S)$ instead of $C_G(S)$ when the meaning is clear. Let $\mathfrak{C}(G)$ denote the set of centralisers of a group G . The relation of inclusion then defines a partial order ' $\leq$ ' on $\mathfrak{C}(G)$. We define the infimum of a pair of elements of $\mathfrak{C}(G)$ as obvious way:

$$C(M_1) \wedge C(M_2) = C(M_1) \cap C(M_2) = C(M_1 \cup M_2).$$

Moreover the supremum $C(M_1) \vee C(M_2)$ of elements $C(M_1)$ and $C(M_2)$ of $\mathfrak{C}(G)$ may be defined to be the intersection of all centralisers containing $C(M_1)$ and $C(M_2)$. Then $C(M_1) \vee C(M_2)$ is minimal among centralisers containing $C(M_1)$ and $C(M_2)$. These definitions make $\mathfrak{C}(G)$ into a lattice, called the *centraliser lattice* of G . This lattice is bounded as it has a greatest element, $G = C(1)$, and a least element, $Z(G)$, the centre of G .

If C and C' are in $\mathfrak{C}(G)$ with C strictly contained in C' we write $C < C'$. If C_i is a centraliser, for $i = 0, \dots, k$, with $C_0 > \dots > C_k$ then we call $C_0, \dots, C_k$ a *centraliser chain* of length k . Infinite descending, ascending and doubly-infinite centraliser chains are defined in the obvious way.

Definition 1.1 (cf. [15], [14]). *If there exists an integer d such that the group G has a centraliser chain of length d and no centraliser chain of length greater than d then G is said to have centraliser dimension $\mathbf{cdim}(G) = d$. If no such integer d exists we define $\mathbf{cdim}(G) = \infty$.*

If $\mathbf{cdim}(G) = d$ then every strictly descending chain of centralisers in $\mathfrak{C}(G)$ from G to $Z(G)$ contains at most d inclusions. This number is usually referred to as the *height* of the lattice; so $\mathbf{cdim}(G)$ is the height of the centraliser lattice of G .

Throughout the remainder of this paper G denotes a (finitely generated free) partially commutative group (unless we explicitly indicate otherwise). The centraliser dimension of partially commutative groups is finite because all partially commutative groups are linear [9] and all linear groups have finite centraliser dimension [14].

The centraliser dimension of a partially commutative group is easy to calculate and depends only on the centralisers of subsets of X . If $Y \subseteq X$ then we call $C(Y)$ a *canonical* centraliser. From the above it follows that the intersection of

two canonical centralisers is again a canonical centraliser. As shown in Section 3 below the supremum, in $\mathfrak{C}(G)$, of two canonical centralisers is also a canonical centraliser. Hence the set $\mathfrak{C}(X; G)$ of canonical centralisers forms a sublattice of $\mathfrak{C}(G)$. We show in Theorem 3.3 that the centraliser dimension of G is equal to the height of the lattice $\mathfrak{C}(X; G)$.

Now let x be a vertex of Γ and let Γ_x denote the graph formed from Γ by removal of the vertex x . Let $G = G(\Gamma)$ and let $G_x = G(\Gamma_x)$, the partially commutative group with graph Γ_x . In Theorem 4.4 we show that $\text{cdim}(G) = \text{cdim}(G_x) + d$, where $d = 0, 1$ or 2 . Moreover the theorem describes exactly how d depends on Γ and x . We conclude the paper by applying Theorem 4.4 to calculate the centraliser dimension of groups in some specific classes of partially commutative groups.

The authors thank Claas Röver for helpful conversations and careful reading of, in particular Theorem 3.3.

2 Preliminaries

First we recall some of the notation and definitions of [8]. If $w \in (X \cup X^{-1})^*$ then we denote by $\alpha(w)$ the set of elements $x \in X$ such that x or x^{-1} occurs in w . By abuse of notation we identify words of $(X \cup X^{-1})^*$ with the elements of G which they represent. Moreover for $u, v \in (X \cup X^{-1})^*$ we use $u = v$ to mean u and v are equal as elements of G . Equality of words u, v in $(X \cup X^{-1})^*$ is denoted by $u \equiv v$. If $g \in G$ and $w \in (X \cup X^{-1})^*$ is a word of minimal length representing g then we say that w is a *minimal* form of g (or just that w is *minimal*).

The Transformation Lemma, [8, Lemma 2.3] asserts that, if u and v are minimal and $u = v$ then we can transform the word u into the word v using only commutation relations from R (that is, without insertion or deletion of any subwords of the form $x^\varepsilon x^{-\varepsilon}$, $x \in X$). From the Transformation Lemma it follows that if u and v are minimal and $u = v$ then $\alpha(u) = \alpha(v)$ and $|u| = |v|$ (where $|w|$ denotes the length of the word w). Therefore, for any element $g \in G$ we may define $\alpha(g) = \alpha(w)$, and the *length* $l(g)$ of g as $l(g) = |w|$, where w is a minimal form of g .

If $g, h \in G$ such that $l(gh) = l(g) + l(h)$ then we write $gh = g \circ h$. It follows that $gh = g \circ h$ if and only if, for all minimal forms u and v of g and h , respectively, uv is a minimal form for gh . Clearly if $w \in (X \cup X^{-1})^*$ is minimal and $w \equiv uv$ then, in G , $w = u \circ v$. If $k = g \circ h$ then we say that g is a *left divisor* of k (and h is a *right divisor* of k). We denote by $D_l(k)$ and $D_r(k)$ the set of left and right divisors of k , respectively. We shall use the following Lemma (cf. [8, Lemma 3.9]).

Lemma 2.1. *If $a \circ b = c \circ d$. Then $a = c_1 \circ d_1$ and $b = c_2 \circ d_2$, where $c = c_1 \circ c_2$, $d = d_1 \circ d_2$ and $[\alpha(c_2), \alpha(d_1)] = 1$.*

Proof. If $l(a) = 0$ then the result holds, trivially. Suppose then that $l(a) > 0$ and that the result holds for all a of shorter length. Then we may write $a = x \circ a'$,

where $x \in X^{\pm 1}$. In this case $x \in D_l(c \circ d)$, so either $x \in D_l(c)$ or $x \in D_l(d)$ and $[x, \alpha(c)] = 1$. Assume first that the latter holds. Then $d = x \circ d'$ and it follows that $a' \circ b = c \circ d'$. As $l(a') < l(a)$ there exist c_1, c_2, d'_1 and d_2 such that $a' = c_1 \circ d'_1$, $b = c_2 \circ d_2$, $c = c_1 \circ c_2$, $d' = d'_1 \circ d_2$ and $[\alpha(c_2), \alpha(d'_1)] = 1$. The result now follows on setting $d_1 = x \circ d'_1$. In the case where $x \in D_l(c)$ a similar, simpler, argument shows that the lemma holds. $\square$

Let u and v be elements of G . A *greatest common left divisor* of u and v is an element $c \in D_l(u) \cap D_l(v)$ such that every element of $D_l(u) \cap D_l(v)$ is a left divisor of c . In [8] it is shown that every pair of elements u, v of G has a unique greatest common left divisor, which we denote by $\gcd_l(u, v)$. This definition extends to n -tuples of elements of G ; we recursively define $\gcd_l(u_1, \dots, u_n)$ to be $\gcd_l(\gcd_l(u_1, \dots, u_{n-1}), u_n)$. It follows from [8] that $\gcd_l(u_1, \dots, u_n)$ is independent of the order of $u_1, \dots, u_n$. The *greatest common right divisor* $\gcd_r(u_1, \dots, u_n)$ of an n -tuple of elements is defined analogously.

A subgroup of G generated by a subset Y of G is called an *canonical parabolic subgroup*. Let $Y \subseteq X$ and denote by $G(Y)$ the partially commutative group given by the presentation with generators Y and relators those $[x_i, x_j] \in R$ such that $x_i, x_j \in Y$. Thus $G(Y)$ has commutation graph the full subgraph of $\Gamma(G)$ generated by Y . It follows from the Transformation Lemma, [8, Lemma 2.5], that $G(Y)$ is the canonical parabolic subgroup of G generated by Y . If $g \in G$ and H is a canonical parabolic subgroup of G then we say that H^g is a *parabolic subgroup*. Thus parabolic subgroups are those with generating set $\{y^g : g \in Y\}$, for some subset Y of X and some $g \in G$.

As in [8], given an element $g \in G$ and a subset Y of X , the set $D_l(g, Y)$ of *parabolic left divisors* of g in $G(Y)$ is defined to be $D_l(g) \cap G(Y)$. A *greatest common left divisor* $\gcd_l(g, Y)$ of g in $G(Y)$ is an element d of $D_l(g, Y)$ such that every element of $D_l(g, Y)$ is a left divisor of d . It follows from [8] that $\gcd_l(g, Y)$ exists and is unique, for all elements $g \in G$ and all canonical parabolic subgroups $G(Y)$ of G .

We say that $w \in (X \cup X^{-1})^*$ is *cyclically minimal* if $l(w) \leq l(w^g)$, for all $g \in G$. If w is minimal and $w \equiv uv$ with u and v minimal then we call $vu \in (X \cup X^{-1})^*$ a *cyclic permutation* of w .

Lemma 2.2. *The following are equivalent.*

- (i) w is cyclically minimal.
- (ii) If $y \in X \cup X^{-1}$ is a left divisor of w then y^{-1} is not a right divisor of w .
- (iii) All cyclic permutations of w are minimal.

Moreover, if w is cyclically minimal then $\alpha(w^g) \supseteq \alpha(w)$, for all $g \in G$.

Proof. The equivalence of (i) and (ii) is Proposition 5.6 of [8]. To see that (i) implies (iii) suppose that w is cyclically minimal and that vu is a cyclic permutation of w . Then $vu = v w v^{-1}$ so $l(vu) \geq l(w) = l(u) + l(v)$. Hence vu is minimal as required. To complete the proof of the equivalence we show that

(iii) implies (ii). Suppose that $y \in X^{\pm 1}$ is a left divisor of w and that y^{-1} is a right divisor of w . Then $w = y \circ w_1 \circ y^{-1}$, so the cyclic permutation $w_1 y^{-1} y$ is not minimal, a contradiction.

For the final statement suppose that w is cyclically minimal and that u is a minimal form for an element of G . The result holds trivially if $u = 1$ and we use induction on the length of u . If $u^{-1} w u$ is minimal the result follows so we may assume that this is not the case. If $u = s \circ t$, where s is non-trivial and commutes with w , then $w^u = w^t$, so the result follows by induction. Therefore we may assume that u has no such left divisor. From the Transformation Lemma (see [8, Lemma 2.5]) it follows that if cancellation occurs in the process of reducing w^u to a minimal form then the first such cancellation must involve a letter from w cancelling with a letter from $u^{\pm 1}$. That is, there exists $y \in X^{\pm 1}$ such that $u = y \circ v$ and either $w = y \circ w_0$ or $w = w_1 \circ y^{-1}$. In this case $w^u = (w_0 \circ y)^v$ or $(y^{-1} \circ w_1)^v$. In both cases the result follows by induction. $\square$

If w is a minimal form of an element $g \in G$ and w is cyclically minimal then we say that g is a cyclically minimal element of G . It follows from Lemma 2.2 that if g is a cyclically minimal element of G then all minimal forms of g are cyclically minimal elements of $(X \cup X^{-1})^*$. For a cyclically minimal element $g \in G$ we define $[g] = \{h \in G \mid h = v u, \text{ for some } u, v \text{ such that } g = u \circ v\}$.

From Proposition 3.9 of [6], if $g \in G$ then there exist $u, w \in (X \cup X^{-1})^*$, with w cyclically minimal, such that $g = u^{-1} \circ w \circ u$. Thus g is cyclically minimal if and only if $u = 1$. Observe that if g is cyclically minimal then $l(g^n) = n l(g)$. Therefore partially commutative groups are torsion free.

In Section 3 the following lemma and its corollary will be useful.

Lemma 2.3. *Let w be a cyclically minimal element of G and let $g \in G$. Then $w^g = u^{-1} \circ v \circ u$ and $g = a \circ b \circ u$, where v is cyclically minimal, $[v] = [w]$, $\alpha(a) \subseteq \alpha(w)$, $w^a = v$ and $b \in C(\alpha(w))$.*

Proof. If $l(g) = 0$ then the result clearly holds. Assume inductively that $l(g) > 1$ and that the lemma holds for all g of shorter length. If $w^g = g^{-1} \circ w \circ g$ then the result holds with $u = g$, $w = v$ and $a = b = 1$. Therefore we may assume that $w^g \neq g^{-1} \circ w \circ g$. From [8, Lemma 2.2] either (i) $\gcd_l(g, w) \neq 1$; or (ii) $\gcd_l(g, w^{-1}) \neq 1$; or (iii) $D_l(g) \cap C(\alpha(w)) \neq 1$.

In case (i) suppose $\gcd_l(g, w) = c \neq 1$. Then $g = c \circ g_1$, $w = c \circ w_1$, $w^g = (w_1 c)^{g_1}$ and $l(g_1) < l(g)$. As w is cyclically minimal Lemma 2.2 implies that $w_1 \circ c$ is also cyclically minimal and by induction $(w_1 c)^{g_1} = u^{-1} \circ v \circ u$, with $[v] = [w_1 \circ c] = [w]$, $g_1 = a' \circ b \circ u$, $\alpha(a') \subseteq \alpha(w_1 c) = \alpha(w)$, $(w_1 c)^{a'} = v$ and $b \in C(\alpha(w))$. Then $g = c \circ a' \circ b \circ u$, $\alpha(c \circ a') \subseteq \alpha(w)$ and $v = w^{c \circ a'}$, so the result holds with $a = c \circ a'$. A similar argument works in case (ii).

Assume then that case (iii) holds. Let $c \in D_l(g) \cap C(\alpha(w))$, $c \neq 1$. Then $g = c \circ g_1$, $l(g_1) < l(g)$ and $w^g = w^{g_1}$. By induction $w^{g_1} = u^{-1} \circ v \circ u$, where v is cyclically minimal, $[v] = [w]$, $g_1 = a \circ b' \circ u$, $\alpha(a) \subseteq \alpha(w)$, $w^a = v$ and $b' \in C(\alpha(w))$. As $c \in C(\alpha(w))$ and $\alpha(a) \subseteq \alpha(w)$, we have $[c, \alpha(a)] = 1$ and so $g = c \circ a \circ b' \circ u = a \circ c \circ b' \circ u$. Therefore the lemma holds with $b = c \circ b'$. $\square$

Corollary 2.4. *Let w, g be (minimal forms of) elements of G and $w = u^{-1} \circ v \circ u$, where v is cyclically minimal. Then there exist minimal forms a, b, c, d_1, d_2 and e such that $g = a \circ b \circ c \circ d_2$, $u = d_1 \circ a^{-1}$, $d = d_1 \circ d_2$, $w^g = d^{-1} \circ e \circ d$, $[e] = [v]$, $e = v^b$, $\alpha(b) \subseteq \alpha(v)$ and $[\alpha(b \circ c), \alpha(d_1)] = [\alpha(c), \alpha(v)] = 1$.*

Proof. We have $w^g = v^{ug}$. Reducing ug to a minimal form we have $u = u_1 \circ g_1^{-1}$, $g = g_1 \circ g_2$ and $ug = u_1 \circ g_2$, for appropriate u_1, g_1, g_2 . Then $w^g = v^{u_1 \circ g_2}$ and using Lemma 2.3 $w^g = d^{-1} \circ e \circ d$, $u_1 \circ g_2 = b \circ c \circ d$, where e is cyclically minimal, $[e] = [v]$, $c \in C(\alpha(v))$, $\alpha(b) \subseteq \alpha(v)$ and $e = v^b$. It follows from Lemma 2.1 that $u_1 = b_1 \circ c_1 \circ d_1$, $g_2 = b_2 \circ c_2 \circ d_2$, where $b = b_1 \circ b_2$, $c = c_1 \circ c_2$, $d = d_1 \circ d_2$ and $[\alpha(b_2), \alpha(c_1 \circ d_1)] = [\alpha(c_2), \alpha(d_1)] = 1$. Then $\alpha(b) \subseteq \alpha(v)$, and $v^b = e \in [v]$ implies that $\alpha(b_i) \subseteq \alpha(v)$ and $v^{b_1} \in [v]$. Also $c \in C(\alpha(v))$ implies $c_i \in C(\alpha(v))$. Now $u = u_1 \circ g_1^{-1} = b_1 \circ c_1 \circ d_1 \circ g_1^{-1}$ and $w = u^{-1} \circ v \circ u$, $v^{b_1} \in [v]$ imply $b_1 = 1$. Then $c_1 \in C(\alpha(v))$ implies $c_1 = 1$. Hence $u = d_1 \circ g_1^{-1}$, $g = g_1 \circ b \circ c \circ d_2$ and the result follows with $g_1 = a$. $\square$

We shall use one of the normal forms for elements of G defined in [8]. In order to describe this we introduce the *non-commutation* graph $\Phi = \Phi(G)$ of G . This is the undirected, simple graph with vertex set X and an edge joining vertices x and y if and only if $[x, y] \notin R$. Also, for $Y \subseteq X$, denote by $\Phi(Y)$ the full subgraph of Φ with vertex set Y ; so $\Phi(Y) = \Phi(G(Y))$. Extend this notation to arbitrary subsets S of G by setting $\Phi(S) = \Phi(\alpha(S))$, where $\alpha(S) = \cup_{s \in S} \alpha(s)$. If $S = \{g\}$, a set with one element, we write $\Phi(g)$ for $\Phi(S)$. Suppose that the connected components of Φ are $\Phi_1, \dots, \Phi_k$ and let X_s be the vertex set of Φ_s ; so $X = \cup_{s=1}^k X_s$. Then $\Phi_s = \Phi(X_s)$ and $G = G(X_1) \times \dots \times G(X_k)$.

Let g be a cyclically minimal element of G . If $\Phi(g)$ is connected then we say that g is a *block*. Otherwise, suppose that $\Phi(g)$ has connected components $\Phi_1, \dots, \Phi_k$. Then g has a minimal form $w = w_1 \cdots w_k$, where w_s is a block with $\Phi(w_s) = \Phi_s$, and consequently $[w_s, w_t] = 1$, for all $s, t \in \{1, \dots, k\}$. We call $w_1 \cdots w_k$ a *block decomposition* of g . Note that it follows from the Transformation Lemma that if g has another block form $u = u_1 \cdots u_l$ then $k = l$ (as $\Phi(u) = \Phi(g) = \Phi(w)$) and after reordering the u_s 's if necessary $w_s = u_s$, for $s = 1, \dots, k$. Now let g be an arbitrary element of G and write $g = u^{-1} \circ w \circ u$, with w cyclically minimal. Let $w_1 \cdots w_n$ be a block decomposition of w . Then $g = w_1^u \cdots w_n^u$ and we say that $v_i = w_i^u$ is a *block* of g and that g has *block decomposition* $v_1 \cdots v_n$. In this case a straightforward induction on $l(u)$ shows that, since $\Phi(w_i)$ is connected, $\Phi(v_i)$ is connected, for all i .

Following [6], given $g \in G$ we define $a \in G$ to be a *root* of g if the following conditions hold.

- (i) $a^m = g$, for some $m \geq 1$ and
- (ii) if $b^n = g$, for some $b \in G$ and $n \geq 1$, then $n|m$.

As shown in [6] every element of G has a unique root, which we denote $r(g)$. If g is its own root then we say that g is a *root element*. Put another way, g is a root element if g is not a proper power of any element of G . (In fact in [6] the definition of root is made in a slightly wider context and also requires

that, in (ii), $b = a^{m/n}$. However it is also shown there that, if G is a partially commutative group and if $g, h \in G$ with $g^n = h^n$ then $g = h$; so our definition is equivalent for partially commutative groups.)

If $w \in G$ define $A(w) = G(Y)$, where Y is the set of elements of $X \setminus \alpha(w)$ which commute with every element of $\alpha(w)$. If $S \subseteq G$ then we define $A(S) = \cap_{w \in S} A(w)$. Let w be a cyclically minimal element of G with block decomposition $w = w_1 \cdots w_k$ and let $v_i = r(w_i)$. Then, from [6, Theorem 3.10],

$$C(w) = \langle v_1 \rangle \times \cdots \times \langle v_k \rangle \times A(w). \quad (2.1)$$

It follows that

$$C(w) = \cap_{i=1}^k C(v_i). \quad (2.2)$$

Furthermore, if $u = z^{-1} \circ w \circ z$ then $C(u) = [C(w)]^z$ so

$$C(u) = \langle v_1^z \rangle \times \cdots \times \langle v_k^z \rangle \times [A(w)]^z, \quad (2.3)$$

but note that in general $[A(w)]^z \neq A(u)$.

We shall also need the following lemma, which is based on Proposition 5.7 of [8], and its corollary.

Lemma 2.5. *Let u and v be cyclically minimal with block decompositions $u = u_1 \cdots u_k$ and $v = v_1 \cdots v_l$. If $u = v^g$, for some $g \in G$ then $k = l$ and, after reordering of the v_s 's if necessary, $u_s = v_s^g$ and some minimal form of u_s is equal to a cyclic permutation of v_s , for $s = 1, \dots, k$.*

Proof. It follows from the proof of Proposition 5.7 of [8] that $k = l$ and that, after an appropriate reindexation, $u_s = v_s^a$, for some $a \in G$ and for all s . Then $u = v^a = v^g$ which implies that $v = v^{ag^{-1}}$, so $ag^{-1} \in C(v)$. This implies that $g = za$, for some $z \in C(v)$, and so, from (2.2), $z \in C(v_s)$, for $s = 1, \dots, k$. Thus $u_s = v_s^a = v_s^g$, for all s , as claimed. The final assertion follows since u_s is minimal. $\square$

Corollary 2.6. *Let u and v be cyclically minimal, root, block elements and let g be such that $u^g = g^{-1} \circ u \circ g$. If u and u^g are in $C(v)$ then $g \in C(v)$.*

Proof. From (2.1) and the fact that u and v are block, root elements and $u \in C(v) = \langle v \rangle \times A(v)$ we have $u \in \langle v \rangle$ or $u \in A(v)$. Thus either $u = v^{\pm 1}$, and without loss of generality $u = v$, or $u \in A(v)$.

Suppose first that $u \notin A(v)$, so $u = v$. As u^g is a block, $\Phi(u^g)$ is connected so $u^g \in C(v)$ implies $u^g \in \langle v \rangle$ or $u^g \in A(v)$. Since u^g is minimal as written, if $u^g \in A(v)$ then $u \in A(v)$, a contradiction. As v is cyclically minimal, if $u^g \in \langle v \rangle$, and the given hypotheses hold, then $g = 1$; so the result follows.

Now suppose that $u \in A(v)$. Then, since it's a block, $g^{-1} \circ u \circ g \in A(v)$, so $g \in A(v)$ and the result follows. $\square$

3 Centraliser dimension and canonical centralisers

In this section we prove that the height of the centraliser lattice $\mathfrak{C}(G)$ of G coincides with the height of a sublattice: namely the lattice of canonical centralisers $\mathfrak{C}(X; G)$. First we must show that $\mathfrak{C}(X; G)$ is indeed a sublattice of $\mathfrak{C}(G)$ and to do so we need to describe the structure of canonical centralisers.

Let Y be a subset of X and define the set of generators *orthogonal* to Y to be $Y^\perp = \{x \in X \mid [x, y] = 1, \text{ for all } y \in Y\}$. Then $G(Y^\perp) \subseteq C(Y)$. On the other hand, if $z \notin G(Y^\perp)$ then there exists $x \in \alpha(z)$ such that $x \notin Y^\perp$. Therefore there is $y \in Y$ such that $[x, y] \neq 1$, that is $x \notin C(y)$. Now from (2.1), $C(y) = \langle y \rangle \times A(y)$, so $x \notin C(y)$ implies $z \notin C(y)$. It follows that $C(Y) = G(Y^\perp)$, a parabolic subgroup. To rephrase this in terms of the notation of Section 2 let $Y_0 = Y \cap Y^\perp = \{y_1, \dots, y_n\}$ and $Z = Y^\perp \setminus Y$, so $Y^\perp = Y_0 \cup Z$. Then $y \in Y_0$ implies $[y, u] = 1$, for all $u \in Y^\perp$, and moreover $G(Z) = A(Y)$. Therefore we have;

$$\text{if } Y \subseteq X \text{ then } C(Y) = G(Y^\perp) = \langle y_1 \rangle \times \dots \times \langle y_n \rangle \times A(Y), \quad (3.1)$$

where $\{y_1, \dots, y_n\} = Y^\perp \cap Y$.

Lemma 3.1. *The set of canonical centralisers $\mathfrak{C}(X; G)$ is a sublattice of $\mathfrak{C}(G)$.*

Proof. It is clear that the intersection of two canonical centralisers is a canonical centraliser. Now let $C_1 = C(Y_1)$ and $C_2 = C(Y_2)$ be canonical centralisers, for some subsets Y_1 and Y_2 of X . Then, in the lattice $\mathfrak{C}(G)$ we have $C_1 \vee C_2 = C$, where C is the intersection of all centralisers containing C_1 and C_2 . Then $C = C(W)$, for some subset $W \subseteq G$, and we set $Z = \cup_{w \in W} \alpha(w)$, so $Z \subseteq X$. Clearly $C(Z) \subseteq C(W)$. From (3.1), $C(Y_1) = G(Y_1^\perp)$. If $a \in Y_1^\perp$ then $a \in C_1 \subseteq C(W)$, so $a \in C(w)$, for all $w \in W$. As $a \in X$ it follows that $a \in C(\alpha(w))$, for all $w \in W$. Therefore $Y_1^\perp \subseteq C(Z)$ and so $C_1 \subseteq C(Z)$. Similarly $C_2 \subseteq C(Z)$ so we have $C = C_1 \vee C_2 \subseteq C(Z) \subseteq C$. Hence $C = C(Z)$, a canonical centraliser. That is $C_1 \vee C_2 \in \mathfrak{C}(X; G)$ and this completes the proof of the lemma. $\square$

Expression (2.1) for the centraliser of cyclically minimal elements gives the structure of the centraliser of certain sets of more than one element as follows.

Lemma 3.2. *Let $w_1, \dots, w_{k-1}, u$ be cyclically minimal root, block elements of G . Let $w_k = g^{-1} \circ u \circ g$, for some $g \in G$ such that either $g = 1$ or $g \notin C(w_1, \dots, w_{k-1})$. Suppose that*

$$C(w_1) > C(w_1, w_2) > \dots > C(w_1, \dots, w_k) \quad (3.2)$$

is a saturated chain of centralisers. Then, there exist integers $m \geq 0$ and $i_1, \dots, i_m$, with $1 \leq i_1 < \dots < i_m \leq k-1$, such that

(i) *if $g = 1$ and $w_k \in C(w_1, \dots, w_{k-1})$ then*

$$C(w_1, \dots, w_k) = \langle w_{i_1} \rangle \times \dots \times \langle w_{i_m} \rangle \times \langle w_k \rangle \times A(w_1, \dots, w_k), \quad (3.3)$$

(ii) and otherwise

$$C(w_1, \dots, w_k) = \langle w_{i_1} \rangle \times \dots \times \langle w_{i_n} \rangle \times (A(w_1, \dots, w_{k-1}) \cap A^g(u)). \quad (3.4)$$

Proof. When $k = 1$ this is (2.1) or (2.3). Write $C_i = C(w_1, \dots, w_i)$ and $A_i = A(w_1, \dots, w_i)$. Assume inductively that

$$C_{k-1} = \langle w_{i_1} \rangle \times \dots \times \langle w_{i_n} \rangle \times A_{k-1},$$

where $n \geq 0$ and $1 \leq i_1 < \dots < i_n \leq k-1$. (Here $w_{i_n} = w_{k-1}$ if and only if $w_{k-1} \in C_{k-2}$.) We have

$$C_k = C_{k-1} \cap (\langle w_k \rangle \times A^g(u)). \quad (3.5)$$

If $g = 1$ and $w_k = u \in A_{k-1}$ then $w_{i_1}, \dots, w_{i_n} \in C(w_k)$. As w_{i_j} is a block this implies that $w_{i_j} \in \langle w_k \rangle$ or $w_{i_j} \in A(w_k)$. If $w_{i_j} \in \langle w_k \rangle$ then $C(w_{i_j}) = C(w_k)$, so (3.2) is not strictly descending, contrary to hypothesis. Hence $w_{i_j} \in A(w_k)$, for $j = 1, \dots, n$. A straightforward check, using 3.5, now shows that

$$C_k = \langle w_{i_1} \rangle \times \dots \times \langle w_{i_n} \rangle \times \langle w_k \rangle \times A_k,$$

as required.

If $g = 1$ and $w_k \notin A_{k-1}$ then $w_k \in C_k$ implies $w_k \in \langle w_i \rangle$, for some i with $1 \leq i \leq k-1$. This implies $C(w_k) = C(w_i)$, a contradiction. Hence we may assume $g \neq 1$ or $w_k \notin C_{k-1}$.

If $g \neq 1$ and $w_k \in A_{k-1}$ then, since u^g is minimal, we have $g \in A_{k-1} \subseteq C_{k-1}$, a contradiction. Suppose then that $w_k \notin A_{k-1}$. Let $z \in C_k$. Then, from 3.5,

$$z = w_{i_1}^{\alpha_1} \dots w_{i_n}^{\alpha_n} \cdot a = w_k^\beta \cdot b^g,$$

where $a \in A_{k-1}$ and $b \in A(u)$. Write $a = a_0^{-1} \circ a_1 \circ a_0$ and $b = b_0^{-1} \circ b_1 \circ b_0$, where a_1 and b_1 are cyclically minimal. Note that we have $a_0, a_1 \in A_{k-1}$ and $b_0, b_1 \in A(u)$. Then

$$\begin{aligned} w_{i_1}^{\alpha_1} \dots w_{i_n}^{\alpha_n} \cdot a_1 &= a_0(w_k^\beta b_1^{b_0 g})a_0^{-1} \\ &= a_0(u^\beta b_1)^{b_0 g}a_0^{-1}, \text{ since } b_0 \in A(u), \\ &= (u^\beta b_1)^{b_0 g a_0^{-1}}. \end{aligned}$$

Now $z_1 = w_{i_1}^{\alpha_1} \dots w_{i_n}^{\alpha_n} \cdot a_1$ and $u^\beta b_1$ are cyclically minimal and, if $\beta \neq 0$ then u^β is a block of $u^\beta b_1$. Therefore, from Lemma 2.5, if $\beta \neq 0$ then $(u^\beta)^{b_0 g a_0^{-1}}$ is a block of z_1 . Blocks of z_1 are either of the form $w_{i_j}^{\alpha_j}$ or are blocks of a_1 . However

$$\begin{aligned} (u^\beta)^{b_0 g a_0^{-1}} &= w_{i_j}^{\alpha_j} \text{ implies} \\ (u^\beta)^{g a_0^{-1}} &= w_{i_j}^{\alpha_j} \text{ implies} \\ (u^\beta)^g &= w_{i_j}^{\alpha_j}, \text{ since } a_0 \in A_{k-1}. \end{aligned} \quad (3.6)$$

If $g \neq 1$ then $(u^\beta)^g$ is not cyclically minimal, and so (3.6) is a contradiction. On the other hand if $g = 1$ then (3.6) implies that $u^\beta = w_{i_j}$ so $w_k = u \in C_k$, again a contradiction. Hence, if $\beta \neq 0$ then $(u^\beta)^{b_0 g a_0^{-1}}$ is a block of a_1 and so belongs to A_{k-1} . This implies that $(u^\beta)^g \in A_{k-1}$ and, since u is cyclically minimal, that $w_k = u^g \in A_{k-1}$, another contradiction. Thus $\beta = 0$ and $z_1 = b_1^{b_0 g a_0^{-1}}$. From Lemma 2.5 again, for $j = 1, \dots, n$, either $\alpha_j = 0$ or

$$w_{i_j}^{\alpha_j} = b_{s_j}^{b_0 g a_0^{-1}} \implies w_{i_j}^{\alpha_j} = b_{s_j}^{b_0 g},$$

for some block b_{s_j} of b_1 . Thus, if $\alpha_j \neq 0$ then $w_{i_j} \in A(u)^g \subseteq C(w_k)$ and, since by definition $w_{i_j} \in C_{k-1}$, we have $w_{i_j} \in C_k$ or $\alpha_j = 0$, for $j = 1, \dots, n$. Also $w_{i_1}^{\alpha_1} \dots w_{i_n}^{\alpha_n} \in A^g(u)$ and so $a \in A_{k-1} \cap A^g(u)$. This means that $z \in T$, where

$$T = \langle w_{j_1} \rangle \times \dots \times \langle w_{j_m} \rangle \times (A_{k-1} \cap A^g(u)),$$

with $\{j_1, \dots, j_m\}$ the subset of $\{1, \dots, k-1\}$ such that $w_{j_l} \in C_k$. Hence $C_k \subseteq T$. The opposite inclusion clearly holds and the result follows. $\square$

The centraliser dimension of a partially commutative group G is by definition the height of its centraliser lattice $\mathfrak{C}(G)$. The main theorem of this paper, which follows, shows that this height is the same as the height of the sublattice consisting of centralisers of canonical generators. This means that the centraliser dimension may be calculated directly from the commutation graph of G .

Theorem 3.3. *Let G be a partially commutative group. Then centraliser dimension of G is equal to the height of the lattice $\mathfrak{C}(X; G)$ of canonical centralisers of G .*

Proof. As pointed out above, G has finite centraliser dimension as it is a linear group. Let d be the centraliser dimension of G . We must show that there exist elements $x_1, \dots, x_d \in X$ such that

$$G > C(x_1) > \dots > C(x_1, \dots, x_d) = Z(G)$$

is a centraliser chain in G . Let

$$G = C(1) > C(w_1) > \dots > C(w_1, \dots, w_d) = Z(G) \quad (3.7)$$

be a maximal centraliser chain in G . For $i = 1, \dots, d$, set $C_i = C(w_1, \dots, w_i)$ and $A_i = A(w_1, \dots, w_i)$.

Step 1. For $i = 1, \dots, d$ we may write $w_i = g_i^{-1} \circ u_i \circ g_i$, with u_i cyclically minimal. We shall show in this step that we may assume that u_i is also a block, root element.

Suppose that the block decomposition of u_j has exactly one block, for $j = 1, \dots, i-1$, but the block decomposition of u_i is $u_i = v_1 \dots v_k$, with $k > 1$. As $C_{i-1} \not\leq C(w_i)$ it follows that $C_{i-1} \not\leq g_i^{-1} C(v_t) g_i$, for some t . As (3.7) is maximal and $g_i^{-1} C(v_t) g_i > C(w_i)$ it follows that $g_i^{-1} C(v_t) g_i \cap C_{i-1} = C(w_i) \cap C_{i-1}$. Thus

we may assume that u_i is a block, for $i = 1, \dots, d$. We may also assume that u_i is a root element, since $C(u) = C(v)$, if v is the root of a block u . Thus

$$C(w_i) = \langle u_i \rangle^{g_i} \times A^{g_i}(u_i).$$

Step 2. Now we shall show that we may in addition assume that w_i is cyclically minimal, for all i . Assume that $g_j = 1$, for $j = 1, \dots, i-1$, but that $g_i \neq 1$, that is, w_i is not cyclically minimal. First consider the case where $D_r(g_i) \cap C_{i-1} \neq 1$, so $g_i = a \circ b$, where $b \in C_{i-1}$. Then, writing $h = b^{-1}$, we have $C_j^h = C_j$, for $j = 1, \dots, i-1$. Hence

$$G > C_1 > \dots > C_{i-1} = C_{i-1}^h > C_i^h > C_d^h = Z(G)$$

is a maximal centraliser chain. Since $C_i^h = C(w_1, \dots, w_{i-1}, u_i^g)$ we may replace w_i with u_i^g , in this case. Therefore we may assume that $D_r(g_i) \cap C_{i-1} = 1$. In particular $g_i \notin C_{i-1}$. Since $w_1, \dots, w_{i-1}$ are cyclically minimal, from Lemma 3.2 we have

$$C_j = \langle w_{j,1} \rangle \times \dots \times \langle w_{j,k_j} \rangle \times A_j,$$

where the $w_{j,i}$ are in $\{w_1, \dots, w_j\}$, for all j with $1 \leq j \leq i-1$. In particular we can write

$$C_{i-1} = \langle w_{i_1} \rangle \times \dots \times \langle w_{i_m} \rangle \times A,$$

where $1 \leq i_1 < \dots < i_m \leq i-1$ and $A = A_{i-1}$. If $w_i \in A$ then, since $u_i^{g_i}$ is minimal we have $g_i \in A \subseteq C_{i-1}$, a contradiction. Hence $w_i \notin A$ and from Lemma 3.2, after reordering $i_1, \dots, i_m$, if necessary, we have

$$C_i = \langle w_{i_1} \rangle \times \dots \times \langle w_{i_k} \rangle \times (A \cap A^{g_i}(u_i)), \quad (3.8)$$

where $w_{i_1}, \dots, w_{i_k} \in A^{g_i}(u_i)$, and $w_{i_{k+1}}, \dots, w_{i_m} \notin A^{g_i}(u_i)$, for some $k \leq m$.

Next we show that $A \cap A^{g_i}(u_i) \subseteq A(u_i) \cap C(g_i)$. Suppose that $h \in A \cap A^{g_i}(u_i)$, so there is $f \in A(u_i)$ such that $h = f^{g_i}$. Write $f = k^{-1} \circ l \circ k$, where l is cyclically minimal. From Corollary 2.4, there are elements a, b, c, d_1, d_2, e , such that $g_i = a \circ b \circ c \circ d_2$, $k = d_1 \circ a^{-1}$, $d = d_1 \circ d_2$, $h = f^{g_i} = d^{-1} \circ e \circ d$, $[e] = [l]$, $e = l^b$, $\alpha(b) \subseteq \alpha(e)$ and $1 = [\alpha(b \circ c), \alpha(d_1)] = [\alpha(c), \alpha(l)]$. As $h \in A$ we have $d_1, d_2 \in A$ and $D_r(g_i) \cap C_{i-1} = 1$, so $d_2 = 1$ and $d = d_1$. Also $f \in A(u_i)$ so $a, d, l \in A(u_i)$ and $\alpha(b) \subseteq \alpha(e)$, $[e] = [l]$, implies $e, b \in A(u_i)$. Since $D_l(g_i) \cap C(u_i) = 1$, this implies that $a = b = 1$ so $g_i = c$, $d = k$, $e = l$ and $1 = [\alpha(g_i), \alpha(d)] = [\alpha(g_i), \alpha(e)]$. Hence $h = f \in A(u_i)$ and $h \in C(g_i)$, so $A \cap A^{g_i}(u_i) \subseteq A(u_i) \cap C(g_i)$, as claimed.

Hence $A \cap A^{g_i}(u_i) \subseteq A \cap A(u_i) \cap C(g_i)$. Conversely if $c \in A \cap A(u_i) \cap C(g_i)$ then $c \in A \cap A^{g_i}(u_i)$. Therefore

$$A \cap A^{g_i}(u_i) = A \cap A(u_i) \cap C(g_i). \quad (3.9)$$

From (3.8) and (3.9) we have

$$C_i = \langle w_{i_1} \rangle \times \dots \times \langle w_{i_k} \rangle \times (A \cap A(u_i) \cap C(g_i)). \quad (3.10)$$

Note that $\langle w_{i_1} \rangle \times \cdots \times \langle w_{i_k} \rangle \subseteq A \cap A^{g_i}(u_i)$ implies that $\langle w_{i_1} \rangle \times \cdots \times \langle w_{i_k} \rangle \subseteq A \cap A(u_i)$. Therefore

$$C_{i-1} \cap C(u_i) \geq \langle w_{i_1} \rangle \times \cdots \times \langle w_{i_k} \rangle \times (\langle u_i \rangle \cap A) \times (A \cap A(u_i)) \geq C_i,$$

from (3.10). If $C_{i-1} \subseteq C(u_i) \cap C(g_i)$ then $C_{i-1} \subseteq C(u_i^{g_i}) = C(w_i)$, which cannot happen as the chain (3.7) is maximal. If $C_{i-1} \not\subseteq C(u_i)$ then $C_{i-1} \cap C(u_i) < C_{i-1}$ and we may replace w_i by u_i , since by maximality of (3.7), $C_i = C_{i-1} \cap C(u_i)$ in this case. Therefore we may assume that $C_{i-1} \subseteq C(u_i)$ but that $C_{i-1} \not\subseteq C(g_i)$. As $C_{i-1} \subseteq C(u_i)$ we have $w_{i_j} \in C(u_i)$, for $j = 1, \dots, k$. Hence $w_i = u_i^{g_i} \in C(w_{i_j})$ and $u_i \in C(w_{i_j})$, for $j = 1, \dots, k$. From Corollary 2.6, $g_i \in C(w_{i_j})$ and so, from (3.10), $C_i \subseteq C(g_i)$. Since (3.7) is maximal and we have $C_{i-1} > C_{i-1} \cap C(g_i) \geq C_i$ it follows that $C_{i-1} \cap C(g_i) = C_i$. Hence we may (temporarily) replace w_i by g_i and, as in Step 1, write $g_i = h_0^{-1} \circ h_1 \circ h_0$, with h_1 cyclically minimal and then replace g_i by $h_0^{-1} h_2 h_0$, where h_2 is the root of a block of h_1 . Thus we have replaced w_i with $h_0^{-1} h_2 h_0$ and $l(h_0^{-1} h_2 h_0) < l(w_i)$. We now repeat this process until, eventually, we replace w_i by a cyclically minimal, root, block element. By induction we may therefore assume that w_i is cyclically minimal, for $i = 1, \dots, d$.

Step 3. Finally we show that we may replace each w_i with an element of X . As $C_d = Z(G)$ we have, using Lemma 3.2, $C_d \leq A(w_1, \dots, w_d)$, otherwise $C_i = C_{i+1}$, for some i . Hence, for each $i = 1, \dots, d$ there exists a minimal integer $j = d(i)$ such that $w_i \notin C_j$. Suppose that for some $i \geq 1$ we have $j = d(i) \geq i+1$ and $l(w_i) > 1$. Let r_i be a cyclically minimal block such that $\alpha(r_i) = \alpha(w_i)$, r_i is a root element and r_i does not commute with w_i : for example a cyclic permutation of w_i . Then

$$C(w_1, \dots, w_{j-1}, r_i) = C_{j-1} \cap A(w_i) \geq C_j.$$

Since (3.7) is maximal and $w_i \in C_{j-1} \setminus A(w_i)$ we have $C_{j-1} \cap A(w_i) = C_j$. Therefore we may replace w_j with r_i . If $j = i+1$ this is all we need do.

Otherwise, if $j > i+1$, define $C_i^{(i+1)} = G$ and, for $t = i+1, \dots, j-1$, define $C_t^{(i+1)} = C(w_{i+1}, \dots, w_t)$ and $C'_t = C_{i-1} \cap C_t^{(i+1)} \cap A(w_i)$. Then $C'_t = C_{i-1} \cap C(w_i, r_i) \cap C_t^{(i+1)}$ is a centraliser of $t+1$ elements of G , for $t = i, \dots, j-1$. Also $C'_{j-1} = C_j$, by definition of r_i , and $C_t = \langle w_i \rangle \times C'_t$, for $t = i, \dots, j-1$. As (3.7) is a saturated centraliser chain so is

$$C'_i > \cdots > C'_{j-1}.$$

Hence

$$G > C_1 > \cdots > C_i > C'_i > \cdots > C'_{j-1} = C_j > \cdots > C_d = Z(G)$$

is a maximal centraliser chain. We may therefore replace the sequence $w_1, \dots, w_d$ with the sequence $w_1, \dots, w_i, r_i, w_{i+1}, \dots, w_{j-1}, w_{j+1}, \dots, w_d$ to give a new chain (3.7). In the new chain $d(i) = i+1$. Note that $w_i \in A_{i-1}$ so that

if $h < i$ then $d(h) \neq i$. Hence, beginning with $i = 1$, we may repeat this process for successive values of i until, for all i , either $l(w_i) = 1$ or $d(i) \leq i + 1$.

Now suppose that $l(w_i) = 1$ and that $j = d(i) \geq i + 1$. Write $w_i = x$ for notational convenience. Let y be a letter of $\alpha(w_j)$ which does not commute with x . (Such a y must exist as $d(i) = j$.) Then $A(y) \geq A(w_j)$ so

$$C_{j-1} > C_{j-1} \cap C(y) \geq C_{j-1} \cap C(w_j) = C_j.$$

By maximality of (3.7) we may replace w_j with y . Repeating this for all i such that $l(w_i) = 1$ and that $j = d(i) \geq i + 1$, we obtain a new sequence $w_1, \dots, w_d$ with the property that either

- (i) $l(w_i) = 1$, or
- (ii) $d(i) = d(i + 1) = i + 1$, in which case we call w_i and w_{i+1} a *matched pair*, or
- (iii) $d(i) \leq i$, $l(w_i) > 1$ and w_i is not one of a matched pair,

for all $i = 1, \dots, d$.

Suppose that i is such that neither (i) or (ii) holds. Then $w_i \notin C_{i-1}$, since this would imply $w_i \in A_{i-1}$ and so, from Lemma 3.2, $d(i) \geq i + 1$. Therefore, using Lemma 3.2 again, $C_i = C_{i-1} \cap A(w_i)$. As $C_i < C_{i-1}$ and $A(w_i) = A(\alpha(w_i))$ there exists $y \in \alpha(w_i)$ such that $A(y) \not\geq C_{i-1}$. Since $A(w_i) \leq A(y)$ we have

$$C_{i-1} > C_{i-1} \cap A(y) \geq C_{i-1} \cap A(w_i) = C_i.$$

Hence we may replace w_i with y . This results in one fewer occurrence of an index i which does not satisfy (i) or (ii). Repeating this process we therefore obtain a sequence $w_1, \dots, w_d$ such that (i) or (ii) holds, for all i .

Consider now i such that $l(w_i) > 1$ and $d(i) = i + 1$. Then as above we may assume $w_{i+1} = r_i$. Choose letters $x, y \in w_i$ which do not commute. These exist as w_i is a block of length more than one. As $\alpha(w_i) \subseteq A_{i-1}$ we have $x, y \in A_{i-1}$, so $x, y \in C_{i-1}$. Define $C'_i = C_{i-1} \cap C(x)$ and $C'_{i+1} = C_{i-1} \cap C(x, y)$. Then $C(x, y) = A(x, y)$ so $C'_{i+1} = C_{i-1} \cap A(x, y) \geq C_{i-1} \cap A(w_i) = C_{i+1}$. As $y \notin C(x)$ and $x \notin C(x, y)$ we have

$$\dots > C_{i-1} > C'_i > C'_{i+1} \geq C_{i+1} > \dots$$

and, by maximality of (3.7), $C'_{i+1} = C_{i+1}$. We may now replace w_i, w_{i+1} by x, y . Making such replacements over all such pairs w_i, w_{i+1} gives the required result. $\square$

4 Extensions of Partially Commutative Groups

In this section we investigate the effect on centraliser dimension of adding or removing a vertex (and its incident edges) from the commutation graph of a

partially commutative group. We shall see that addition of a vertex can either leave the centraliser dimension unaltered or increase it by one or two.

We call a sequence of generators $x_1, \dots, x_l$ of a partially commutative group G a *parameter system* for G if

$$G = C_0 > C_1 > \dots > C_n \supseteq Z(G),$$

where $C_i = C(x_1, \dots, x_i)$ is a centraliser chain in G . We say that the parameter system is *maximal* or *saturated* if the centraliser chain is maximal or saturated, respectively. If $C_n = Z(G)$ then we say that the parameter system *ends* in $Z(G)$.

Let G be a partially commutative group with commutation graph Γ and presentation $\langle X | R \rangle$, as before, and fix $x \in X$. Let $X_x = X \setminus \{x\}$, let Γ_x be the graph obtained from Γ by deleting x and $R_x = \{r \in R : r = [a, b], a \neq x, b \neq x\}$. Let G_x be the partially commutative group with commutation graph Γ_x ; so G_x has presentation $\langle X_x | R_x \rangle$. Let $Y = Y_x = \{y \in X_x : [y, x] = 1 \text{ in } G\}$, let $W = W_x = X_x \setminus Y$ and, as usual, let $A = A_G(x) = G(Y)$; so A is a canonical parabolic subgroup of both G and G_x . Since the above presentation for G can be written as

$$\langle X_x, x \mid R_x, [x, y] = 1 \text{ for } y \in Y \rangle$$

it is easy to see that G is the *HNN*-extension $\langle G_x, x \mid x^{-1}ax = a, \forall a \in A \rangle$.

Now suppose that $P = x_1, \dots, x_n$ is a parameter system for G_x and let $C_i = C_{G_x}(x_1, \dots, x_i)$. In order to understand the effect on centraliser dimension of forming the *HNN*-extension G from G_x we need the following definitions. If there is some l such that $\{x_1, \dots, x_l\} \subseteq Y$ and some $w \in W$ such that $C_{G_x}(w) \supseteq C_l$ then we say that P is *locked* at l with key w (with respect to A). As we shall see below, if P is locked at l with key w then the sequence $x_1, \dots, x_l, w, x_{l+1}, \dots, x_n$ is a parameter system for G .

Now let k be maximal such that $C_k \not\subseteq A$. We say that P is *tied* at k , of type T1a, T1b or T2, at k if either

- T1. a. $k < n$ and there exists $y \in Y$ such that $y \in C_k \setminus C_{k+1}$; or
- b. $k = n$ and there exists $w \in W \cap C_n$; or
- T2. $k < n$, $\{x_1, \dots, x_k\} \subseteq Y$ and $x_{k+1} \in W$.

Again, if P is tied at k then it turns out that $x_1, \dots, x_k, x, x_{k+1}, \dots, x_n$ is a parameter system for G .

Given a maximal parameter system P for G_x we shall see that P is a maximal parameter system for G if and only if P is neither locked nor tied. Moreover we shall describe below how any maximal parameter system Q for G induces a parameter system for G and see that if this parameter system is shorter than Q then it is either locked or tied. As a result it will follow (see Theorem 4.4) that the centraliser dimension of G is at most 2 more than that of G_x .

Before using these definitions to state our theorem we illustrate them with some examples.

Example 4.1. In each of the following cases G is an HNN -extension of G_x with stable letter x , as above.

1. In the group with commutation graph Γ_1 of Figure 1 we have $Y = \{a, b, c\}$ and $W = \{d, e, f\}$. By inspection $\text{cdim}(G) = 5$ and $\text{cdim}(G_x) = 4$. Among the maximal parameter systems of G_x are the following.

- (i) d, f, a, c which is tied at 2 ($x_2 = f$), of type T1a, but not locked. The corresponding centraliser chain is

$$G_x > \langle d, a, b, c \rangle > \langle a, c \rangle > \langle a \rangle > 1.$$

- (ii) c, a, b, f with centraliser chain

$$G_x > \langle f, d, e \rangle > \langle f, d \rangle > \langle d \rangle > 1.$$

This is tied of type T2 at 3 ($x_3 = b$) and is also locked at 3 with key d .

- (iii) c, a, f, b which is neither locked nor tied. The corresponding centraliser chain is

$$G_x > \langle f, d, e \rangle > \langle f, d \rangle > \langle f \rangle > 1.$$

2. In the group with commutation graph Γ_2 of Figure 1 we have $Y = \{a, b, c\}$ and $W = \{d, e\}$. In this case $\text{cdim}(G) = 6$ and $\text{cdim}(G_x) = 4$. G_x has a maximal parameter system b, a, c, e with centraliser chain

$$G_x > \langle a, b, c, d \rangle > \langle a, b, d \rangle > \langle b, d \rangle > \langle d \rangle.$$

This is locked at 3 ($x_3 = c$), with key d , and tied at 4 ($x_4 = e$), of type T1b.

3. In the group with commutation graph Γ_3 of Figure 1 we have $Y = \{c, d\}$ and $W = \{a, b\}$. Here $\text{cdim}(G) = \text{cdim}(G_x) = 4$. G_x has a maximal parameter system c, b, d, a with centraliser chain

$$G_x > \langle a, b, c \rangle > \langle b, c \rangle > \langle b \rangle > 1.$$

This is neither locked nor tied.

4. In the group with commutation graph Γ_4 of Figure 2 we have $Y = \{a, b, c\}$ and $W = \{d, e\}$. Here $\text{cdim}(G) = 6$ and $\text{cdim}(G_x) = 4$. G_x has a maximal parameter system b, a, c, e with centraliser chain

$$G_x > \langle a, b, c, d \rangle > \langle a, b, d \rangle > \langle b, d \rangle > 1.$$

This is locked at 3 ($x_3 = c$), with key d , and tied at 3, of type T1a.




Figure 1: Example 4.1

5. In the group with commutation graph Γ_5 of Figure 2 we have $Y = \{a, b, c\}$ and $W = \{d, e, f, g, h, i, j, k\}$. Here $\text{cdim}(G_x) = 5$, since its commutation graph has degree 3 and l, i, k, j, h is a parameter system for G_x . From Example 4.1.4 above G must have a parameter system of length 6 and as Γ_5 has degree 4 it follows that $\text{cdim}(G) = 6$. We remark that from Example 4.1.4 G_x has parameter system of length 4 which is both locked and tied of type T1.

Let $P = v_1, \dots, v_m$ be a maximal parameter system for G . Let $D_s = C_G(v_1, \dots, v_s)$ and let $D_0 = G$. If x occurs in P with $x = v_j$ then define $j_x = j$. Otherwise set $j_x = m + 1$. If $x \notin Z(G)$ then define i_x to be the minimal integer $i > 0$ such that $v_i \in W$. Such $i \leq m$ exists as $x \notin Z(G) = D_m$; so there must be some element of W in P . If $x \in Z(G)$ then set $i_x = m + 1$. Note that in this case $W = \emptyset$ and x does not occur in P , so that $j_x = m + 1$. Suppose now that x occurs in P and $j_x = j$. Then $D_{j-1} \not\subseteq C_G(x)$. In this case suppose that $D_{j-1} \cap C_G(v_{j+1}) \not\subseteq C_G(x)$. As P is a parameter system $D_{j-1} > D_{j-1} \cap C_G(v_{j+1})$. There exists $u \in X$ such that $u \in (D_{j-1} \cap C_G(v_{j+1})) \setminus C_G(x)$, so $u \neq x$ and we have $D_{j-1} \cap C_G(v_{j+1}) > D_{j-1} \cap C_G(v_{j+1}, x) = D_{j+1}$. Hence we may replace P with the parameter system $v_1, \dots, v_{j-1}, v_{j+1}, x, v_{j+2}, \dots, v_d$. Continuing this way we eventually replace P with a parameter system such that

$$D_{j_x-1} \cap C_G(v_{j_x+1}) \subseteq C_G(x). \quad (4.1)$$

Thus, if x occurs in P then G has a maximal parameter system with this property. Assume now that x occurs in P and that (4.1) holds. Let $j_x = j$ and $i_x = i$. Suppose that $i = j + 1$ and there is $z \in W$ such that $C_{G_x}(z) \supseteq C_{G_x}(v_1, \dots, v_{j-1})$. Then we replace x by z in P . This is possible since $C_{G_x}(z) \supseteq C_{G_x}(v_1, \dots, v_{j-1})$ implies that the only element of $X \cap D_{j-1}$




Figure 2: Example 4.1

not in $C_G(z)$ is x . As $D_{j-1} \cap C_G(v_{j+1}) \subseteq C_G(x)$ and $D_{j-1} \not\subseteq C_G(v_{j+1})$ there must be $w \in W$ such that $w \in D_{j-1}$; so $w \in D_{j-1} \cap C_G(z)$ and $w \notin C_G(v_{j+1})$. Hence after replacing x with z we still have a maximal parameter system for G . In this case we may replace P with a parameter system in which x does not occur. Thus we may assume that if P contains x and $i_x = j_x + 1$ then there is no $z \in W$ such that $C_{G_x}(z) \supseteq C_{G_x}(v_1, \dots, v_{j_x-1})$. We call a parameter system which satisfies (4.1) and the property that either $i_x \neq j_x + 1$ or

$$C_{G_x}(z) \not\supseteq C_{G_x}(v_1, \dots, v_{j_x-1}), \text{ for all } z \in W,$$

a *good* parameter system.

Assume now that P is a good maximal parameter system for G with $i_x = i$ and $j_x = j$. Then either $x \in Z(G)$ or $x \in D_{i-1} \setminus C_G(v_i)$. If either

E1. $x \in Z(G)$ or

E2. $i = j + 1$ or

E3. $i \neq j + 1$ and $D_{i-1} \setminus C_G(v_i)$ contains $u \in X_x$,

then we say that P *extends over W (at i)*. Now define integers k, l and d and a sequence $R = R(P) = y_1, \dots, y_d$ associated to P consisting of elements of X_x as follows.

- (i) $R = v_1, \dots, v_{j-1}, v_{j+1}, \dots, v_{i-1}, v_{i+1}, \dots, v_d$, $d = m - 2$, $k = j - 1$ and $l = i - 2$ if x occurs in P , P does not extend over W and $i > j$.
- (ii) $R = v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_{j-1}, v_{j+1}, \dots, v_d$, $d = m - 2$, $k = j - 2$ and $l = i - 1$ if x occurs in P , P does not extend over W and $i < j$.

- (iii) $R = v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_d$, $d = m - 1$, and $l = i - 1$ if x does not occur in P and P does not extend over W .
- (iv) $R = v_1, \dots, v_{j-1}, v_{j+1}, \dots, v_d$, $d = m - 1$, and $k = j - 1$ if x occurs in P and P extends over W .
- (v) $R = v_1, \dots, v_d$, $d = m$, if x does not occur in P and P extends over W .

Lemma 4.2. *Let P be a good maximal parameter system for G . Then $R = R(P)$ is a parameter system of length d for G_x , where $0 \leq \text{cdim}(G) - d \leq 2$. Moreover exactly one of the following conditions R1–R4 holds.*

- R1. R is locked and tied, of type T1 but not T2, $d = m - 2$, x occurs in P and P does not extend over W .
- R2. R is locked $d = m - 1$, x does not occur in P and P does not extend over W .
- R3. R is tied, $d = m - 1$, x occurs in P and P extends over W .
- R4. $d = m$, x does not occur in P and P extends over W .

Proof. First we show that R is a parameter system for G_x . If $1 \leq s \leq m$ and $s \neq i$ then there exists $u \in X_x$ such that $u \in D_{s-1} \setminus D_s$. Therefore in cases (i) to (iii) R is a parameter system for G_x . If P extends over W at i , $x \notin Z(G)$ and $i \neq j + 1$ then there exists $u \in X_x$ such that $u \in D_{i-1} \setminus D_i$. Hence the required result holds in case (iv) when $i \neq j + 1$. Suppose that P extends over W , $x \notin Z(G)$ and $i = j + 1$. Then $x = v_j = v_{i-1}$ so $C_G(v_1, \dots, v_{i-2}) \not\subseteq C_G(x)$ and $C_G(v_1, \dots, v_{i-2}) \cap C_G(v_i) \subseteq C_G(x)$. Hence there exists $w \in X \setminus C_G(x)$ such that $w \in C_G(v_1, \dots, v_{i-2})$, with $w \notin C_G(v_i)$. Thus $w \in X_x$ and $C_{G_x}(v_1, \dots, v_{i-2}) > C_{G_x}(v_1, \dots, v_{i-2}) \cap C_{G_x}(v_i)$. It follows that R is a parameter system for G_x in case (iv). In case (v) if $x \in Z(G)$ then $W = \emptyset$ so $P = R$ is a parameter system for G_x . Otherwise, in case (v), $i \neq j + 1$ and the same arguments as in case (iv) apply.

Clearly $0 \leq \text{cdim}(G) - d \leq 2$ so it remains to prove that exactly one of R1 to R4 holds. If P does not extend over W then $i \neq j + 1$ and

$$D_{i-1} \setminus C_G(v_i) \cap X = \{x\}. \quad (4.2)$$

In this case suppose first that $i < j$ (so we are in case (ii) or (iii)). Then $v_i \in W$, $l = i - 1$ and $x \notin \{v_1, \dots, v_{i-1}\} \subseteq Y$ and (4.2) implies $C_{G_x}(y_1, \dots, y_l) \subseteq C_{G_x}(v_i)$. Thus R is locked at l with key v_i .

Now suppose that P does not extend over W and that $i > j$ (so we have case (i)). This time $D_{i-1} = C_G(v_1, \dots, x, \dots, v_{i-1})$ and (4.1) implies that $C_G(v_1, \dots, v_{j-1}, v_{j+1}) \subseteq C_G(x)$. Thus (4.2) reads

$$[C_G(v_1, \dots, v_{j-1}, v_{j+1}, \dots, v_{i-1}) \cap C_G(x)] \setminus C_G(v_i) \cap X = \{x\}$$

so that

$$[C_G(v_1, \dots, v_{j-1}, v_{j+1}, \dots, v_{i-1})] \setminus C_G(v_i) \cap X = \{x\}$$

which implies $C_{G_x}(y_1, \dots, y_l) \subseteq G_x(v_i)$. Again R is locked at l with key v_i . Therefore, if P does not extend over W we have a locked parameter system R for G_x . The length of R is $d = \text{cdim}(G) - 2$, if x occurs in P , and $d = \text{cdim}(G) - 1$ otherwise. On the other hand, considering cases (i)-(v), if P extends over W then R has length $d \geq \text{cdim}(G) - 1$.

Now suppose that x occurs in P . Then $x = v_j$ and $D_{j-1} \not\subseteq C_G(x)$ so

$$C_{G_x}(y_1, \dots, y_k) \not\subseteq A. \quad (4.3)$$

Firstly suppose that $i \neq j + 1$ and $j \neq m$. We claim that in this case

$$C_{G_x}(y_1, \dots, y_k) \cap C_{G_x}(y_{k+1}) \subseteq A. \quad (4.4)$$

To see this note that if $i > j$ or P extends over W then $\{y_1, \dots, y_k\} = \{v_1, \dots, v_{j-1}\}$ (see cases (i) and (iv)). Therefore in these cases, as $y_{k+1} = v_{j+1}$, (4.1) implies that (4.4) holds. If P does not extend over W and $i < j$ then $k > l$ and $\{y_1, \dots, y_l\} = \{v_1, \dots, v_{i-1}\}$. Thus if $u \in X \cap C_{G_x}(y_1, \dots, y_k)$ and $u \notin C_G(v_i)$ then $u \in D_{i-1} \setminus C_G(v_i)$, so P extends over W , a contradiction. Hence such u must belong to $C_G(v_i)$ and therefore to D_{j-1} . Now (4.1) implies that if in addition $u \in C_{G_x}(y_{k+1})$ then $u \in C_G(x)$ so $u \in A$. Hence (4.4) holds in this case as well. As $D_j \subseteq C_G(x)$ and $D_j > D_{j+1}$ there exists $y \in Y \cup \{x\}$ such that $y \in D_j$ and $y \notin D_{j+1}$. As $i = j + 1$ we have $y \neq x$, so $y \in Y$ and therefore $y \in C_{G_x}(y_1, \dots, y_k) \setminus C_{G_x}(y_{k+1})$. As $i \neq j + 1$ and $j \neq m$ we have $k < n$ and $y_{k+1} \notin W$. Hence R is tied at k of type T1a (and not of type T1b or T2).

Secondly suppose that $j \neq m$, $i = j + 1$. Then P extends over i , $k = j - 1$ and $y_1, \dots, y_k = v_1, \dots, v_{j-1} \in Y$ (case (iv)). As before (4.3) holds and (4.1) implies (4.4). This time $y_{k+1} = v_{j+1} \in W$ so R is tied at k of type T2 (and possibly also of type T1a). Since P is good R is not locked.

Thirdly suppose that $j = m$; in which case $i \neq j + 1$. Then $d = k$ and there exists $z \in X$ such that $z \in C_G(v_1, \dots, v_{m-1})$ but $z \notin C_G(x)$. Hence $z \in W$ and $z \in C_{G_x}(y_1, \dots, y_d)$. Therefore R is tied at k of type T1b (but not T2).

Finally note that R4 occurs if and only if we have case (v). If P does not extend over W and x occurs in P then, from the above, R is locked at l and tied at k of type T1a or T1b, but not T2. In this case R1 is satisfied. If P does not extend over W and x does not occur in P then R is locked at l and R2 holds. If P extends over W and x occurs in P then R3 holds. $\square$

Let $Q = x_1, \dots, x_n$ be a parameter system for G_x . Let $C_s = C_{G_x}(x_1, \dots, x_s)$, and $C_0 = G_x$. Suppose that Q is locked at l . Note that if $l < l' \leq n$ and $x_i \in Y$, for $i = 1, \dots, l'$, then Q is also locked at l' . If Q is locked at l and l is maximal such that $x_i \in Y$, for $i \leq l$, then we say that Q is *right* locked at l . We may always assume that if Q is locked at l then it is right locked at l . Note also that if Q is tied of type T2 and Q is right locked at l then $x_{l+1} \in W$ so that $k = l < n$. Conversely, if Q is both right locked and tied at $l < n$ then it is tied of type T2 (and possibly also of type T1).

Now suppose that Q is either right locked at l with key w or tied at k , or both. Define integers d , i and j and a sequence $S = S(Q) = u_1, \dots, u_d$ of elements of G as follows.

(i) $S = x_1, \dots, x_k, x, x_{k+1}, \dots, x_l, w, x_{l+1}, \dots, x_n$, $d = n + 2$, $i = l + 2$ and $j = k + 1$, if either

I. Q is locked and tied of type T1 and $k < l$; or

II. Q is locked and tied of type T1b and $k = l = n$.

(ii) $S = x_1, \dots, x_l, w, x_{l+1}, \dots, x_k, x, x_{k+1}, \dots, x_n$, $d = n + 2$, $i = l + 1$ and $j = k + 2$, if Q is locked and tied of type T1 and $k > l$.

(iii) $S = x_1, \dots, x_l, w, x_{l+1}, \dots, x_n$, $d = n + 1$, $i = l + 1$, if Q is locked but not tied.

(iv) $S = x_1, \dots, x_k, x, x_{k+1}, \dots, x_n$, $d = n + 1$, $j = k + 1$, if either Q is not locked but is tied of type T1 or Q is tied of type T2.

(v) $S = x_1, \dots, x_n$, $d = n$, if Q is neither locked nor tied.

Lemma 4.3. *Let Q be a parameter system for G_x . Then $S(Q)$ is a parameter system for G .*

Proof. Let $E_s = C_G(u_1, \dots, u_s)$, for $s = 1, \dots, d$, and $E_0 = G$. First suppose that Q is locked and tied of type T1 and $k < l$. Clearly $E_0 > E_1 > \dots > E_{j-1}$. As Q is tied at k with $k < l \leq n$ it must be tied of type T1a, so k is maximal such that $C_k \not\subseteq A$ and there exists $y \in Y$ and $z \in W$ such that $y, z \in C_k$ but $y, z \notin C_{k+1}$. Hence $y, z \in E_{j-1}$, $y \in E_j$, $z \notin E_j$ and $y \notin E_{j+1}$; so $E_{j-1} > E_j > E_{j+1}$. Since $C_{k+1} \subseteq A$ we have $C_G(x_1, \dots, x_k) \cap C_G(x) \cap C_G(x_{k+1}, \dots, x_t) = C_G(x_1, \dots, x_k) \cap C_G(x_{k+1}, \dots, x_t)$, for $t = k + 1, \dots, i - 1$, so we also have $E_{j+1} > \dots > E_{i-1}$. As Q is locked at l with key w we have $x_i \in Y$, for $i = 1, \dots, k, k + 1, \dots, l$, and $C_{G_x}(w) \supseteq C_l$. Hence $x \in C_G(u_1, \dots, u_{i-1})$ and $x \notin C_G(u_1, \dots, u_{i-1}, w)$. Moreover, if $z \in X$ and $z \in C_G(u_1, \dots, u_{i-1}) \setminus C_G(w)$ then $z = x$. As Q is right locked at l , $u_{i+1} = x_{l+1} \in W$ so $C_G(u_1, \dots, u_{i-1}, w, u_{i+1}, \dots, u_t) = C_G(u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_t)$, for all $t \geq i + 1$. It follows that $E_{i-1} > E_i > E_{i+1} > \dots > E_d$ and so S is a parameter system for G .

If Q is locked and tied of type T1b and $k = l = n$ then $E_0 > \dots > E_n$. As $x, w \in E_n$, $x \in E_{n+1} \setminus E_{n+2}$ and $w \notin E_{n+1}$ it follows that $S(Q)$ is a parameter system for G .

Now suppose that Q is locked and tied of type T1 and $k > l$. As above $E_1 > \dots > E_{i-1} > E_i > E_{i+1} > \dots > E_{j-1}$ and $E_{j-1} = C_G(x_1, \dots, x_l) \cap C_G(x_{l+1}, \dots, x_k)$. First assume that $k \neq n$, so that Q is tied of type T1a. By definition of tied there is $y \in Y$ and $z \in W$ such that $\{y, z\} \subseteq C_k \setminus C_{k+1}$ and, from the above, $\{y, z\} \subseteq E_{j-1}$. As in the previous case it now follows that S is a parameter system for G . In the case where $k = n$ we have $E_1 > \dots > E_{d-1}$ and there exists $z \in W$ such that $z \in C_n = Z(G_x)$. Hence $z \in E_{d-1}$ but $z \notin E_d$. Again S is a parameter system for G .

Similar arguments show that in case (iii) and, in the instances of case (iv) when Q is tied of type T1, S is a parameter system for G . Obviously S is a parameter system for G in case (v). This leaves the case where Q is tied

of type T2 at k . Note that if also Q is right locked at l then $k = l$. Then $E_1 > \dots > E_{j-1}$ and there exists $z \in W$ such that $z \in C_k$ but $z \notin C_{k+1}$. Since $C_{k+1} \subseteq A$ we have $C_G(x_1, \dots, x_{k+1}) \subseteq C_G(x)$ and $z \in W$ implies $z \notin C_G(x)$. As $x_s \in Y$, for $s = 1, \dots, k$ we have $x \in C_G(x_1, \dots, x_k)$. Hence $z, x \in E_{j-1}$, $x \in E_j$, $z \notin E_j$ and $x \notin E_{j+1}$, as $x_{k+1} \in W$, so $E_{j-1} > E_j > E_{j+1}$. As before $E_{j+1} > \dots > E_d$, so S is a parameter system for G . Thus in all cases S is a parameter system for G . $\square$

If a parameter system for G_x is tied of type T1 but is not tied of type T2 then we say that it is of type T1-exclusive.

Theorem 4.4. *In the notation above $0 \leq \text{cdim}(G) - \text{cdim}(G_x) \leq 2$ and, more precisely, the following hold.*

1. $\text{cdim}(G) - \text{cdim}(G_x) = 2$ if and only if G_x has a maximal parameter system which is both locked and tied of type T1-exclusive.
2. $\text{cdim}(G) - \text{cdim}(G_x) = 1$ if and only if G_x has a parameter system which is either
 - (a) maximal and locked or tied but not both locked and tied of type T1-exclusive; or
 - (b) of length $\text{cdim}(G_x) - 1$, ends in $Z(G_x)$, and is both locked and tied of type T1-exclusive.
3. Otherwise $\text{cdim}(G) = \text{cdim}(G_x)$.

Proof. As G_x is a subgroup of G it follows that $\text{cdim}(G_x) \leq \text{cdim}(G)$. Suppose that G has a maximal parameter system Q which is locked at l and tied, of type T1-exclusive, at k . As above, $l \neq k$. From Lemma 4.3 $S(Q)$ is a parameter system for G so $\text{cdim}(G) \geq \text{cdim}(G_x) + 2$ and, from Lemma 4.2, we conclude that $\text{cdim}(G) = \text{cdim}(G_x) + 2$.

Conversely, suppose that $\text{cdim}(G) = \text{cdim}(G_x) + 2$ and let P be a maximal parameter system for G . We may assume that P is good. Then if $R = R(P)$ satisfies R2, R3 or R4 we have $\text{cdim}(G) \leq \text{cdim}(G_x) + 1$, a contradiction. Hence R satisfies R1 and since R has length $\text{cdim}(G) - 2 = \text{cdim}(G_x)$ it must be maximal. This proves the first numbered statement of the Theorem.

Now suppose that $\text{cdim}(G) \leq \text{cdim}(G_x) + 1$; so, from Lemma 4.3 G_x has no maximal parameter system which is both locked and tied of type T1-exclusive. Suppose that G_x has a maximal parameter system Q which is either locked or tied. From Lemma 4.3, $S(Q)$ is a parameter system for G of length $\text{cdim}(G_x) + 1$. Hence, in this case $\text{cdim}(G) = \text{cdim}(G_x) + 1$. If G_x has a parameter system Q of length $\text{cdim}(G_x) - 1$ which is both locked and tied of type T1-exclusive then $S(Q)$ has length $\text{cdim}(G_x) + 1$, so again $\text{cdim}(G) = \text{cdim}(G_x) + 1$.

Conversely suppose that $\text{cdim}(G) = \text{cdim}(G_x) + 1$, let P be a good maximal parameter system for G and let $R = R(P)$. Then R does not satisfy R4, since this would imply $\text{cdim}(G_x) \geq \text{cdim}(G)$. From Lemma 4.2, R satisfies one of R1, R2 or R3. If R satisfies R2 or R3 then R has length $\text{cdim}(G) - 1 = \text{cdim}(G_x)$

so is maximal. In this case G_x has a maximal parameter system which is either locked or tied. If R satisfies R1 then R has length $\text{cdim}(G) - 2 = \text{cdim}(G_x) - 1$. If R does not end in $Z(G_x)$ then there exists $u \in X_x$ such that $R' = y_1, \dots, y_d, u$ is a parameter system for G_x , which must be maximal. Since R is locked at $l \leq d$, R' is also locked at l and so G_x has a maximal parameter system which is locked. Thus we may assume that R ends in $Z(G_x)$. This proves the second numbered statement of the theorem and the third follows from Lemma 4.2. $\square$

In some cases we have simple criteria which determine the relationship of $\text{cdim}(G)$ to $\text{cdim}(G_x)$. We sum these up in the next two corollaries. Observe though that, of the extensions of Example 4.1, only 2 falls within the scope of Corollary 4.5.

Corollary 4.5. *In the notation above let $B = G(W)$.*

1. $\text{cdim}(G) = \text{cdim}(G_x) + 2$ if
 - (a) $Z(G) < Z(G_x)$; or
 - (b) $\text{cdim}(A) = \text{cdim}(G_x)$.
2. $\text{cdim}(G) = \text{cdim}(G_x)$ if
 - (a) $G_x = A \times B$ and $Z(B) = 1$; or
 - (b) $C_G(x) = C_G(S)$, for some $S \subseteq G_x$.

Proof. Let $Q = x_1, \dots, x_n$ be parameter system for G_x and let $C_s = C_{G_x}(x_1, \dots, x_s)$.

1. (a) If $Z(G) < Z(G_x)$ then there exists $z \in W$ such that $z \in Z(G_x)$. Suppose that Q is a maximal parameter system for G_x . Then Q is tied of type T1b at n , so is tied of type T1-exclusive. Let l be maximal such that $x_s \in Y$, for all $s \leq l$. Since $z \in Z(G_x)$ we have $C_{G_x}(z) \supseteq C_l$. Hence Q is locked at l and the result follows from Theorem 4.4.
- (b) We may now assume that $Z(G) = Z(G_x)$. Removing all generators in the centre of G results in a new graph group with the same centraliser dimension as G . Therefore we may assume that $Z(G) = Z(G_x) = 1$. Let $T = t_1, \dots, t_d$ be a maximal parameter system for A . This is also a parameter system for G_x , which must be maximal since $\text{cdim}(A) = \text{cdim}(G_x)$. Let $F_s = C_{G_x}(t_1, \dots, t_s)$, for all s . Then $F_d = Z(G_x) = 1$ and so there exists $w \in W$ such that $C_{G_x}(w) \supseteq F_d$. As $t_i \in A$ we have $t_i \in Y$, for all i , so as a parameter system for G_x , T is locked at d with key w . As T is a parameter system for A there is $y_s \in Y$ such that $y_s \in F_{s-1} \setminus F_s$, for $s = 1, \dots, d$. Let k be maximal such that $F_k \not\subseteq A$. As $Z(G_x) = 1$, we have $k < d$. Then T is tied of type T1a at k . From Theorem 4.4 then $\text{cdim}(G) = \text{cdim}(G_x) + 2$.

2. (a) Suppose that Q is a parameter system for G_x which ends in $Z(G_x)$. Let l be maximal such that $x_s \in Y$, for $s = 1, \dots, l$. Then $W \subseteq C_l$ and so if there is $w \in W$ such that $C_{G_x}(w) \supseteq C_l$ then $C_{G_x}(w) \supseteq W$, contradicting the hypothesis that $Z(B) = 1$. Hence Q is not locked. Let k be maximal such that $C_k \not\subseteq A$. Then, as $Z(G_x) = 1$ we have $k < n$. Every element of W commutes with every element of Y , so $C_{k+1} \subseteq A$ implies $x_{k+1} \in W$. Therefore $C_{G_x}(x_{k+1}) \supseteq Y$ and Q cannot be tied of type T1. If Q is tied of type T2 then $x_s \in Y$, for $s = 1, \dots, k$, in which case $B \subseteq C_k$. Therefore $x_{k+1} \in C_{k+1}$, a contradiction. Hence G_x has no parameter system which is either locked or tied and ends in $Z(G)$. From Theorem 4.4, $\text{cdim}(G) = \text{cdim}(G_x)$.
- (b) Since $C_G(x) = C_G(S)$ we have $A = C_{G_x}(S)$, so that $y \in Y$ implies $S \subseteq C_{G_x}(y)$. If $x_t \in Y$, for $t = 1, \dots, l$, then $S \subseteq C_l$. Moreover, if $w \in W$ then $[w, s] \neq 1$, for some $s \in S$, so $S \not\subseteq C_{G_x}(w)$. Therefore Q is not locked. Let k be maximal such that $C_k \not\subseteq A$. Then $C_{k+1} \subseteq A = C_{G_x}(S)$. Thus there is at least one $w \in W$ such that $w \in C_k \setminus C_{k+1}$. Fix such a w . Then $w \notin A$ so there exists some $u \in X_x$ such that $u \in \alpha(S)$ and $[u, w] \neq 1$. As $C_{G_x}(u) \supseteq C_{G_x}(S)$ we have $C_k > C_k \cap C_{G_x}(u) \supseteq C_k \cap C_{G_x}(S) > C_{k+1}$. Hence we may assume that $x_{k+1} = u$. Now if $y \in Y$ then $y \in C_{G_x}(S)$ implies $y \in C_{G_x}(u)$ so that $y \notin C_k \setminus C_{k+1}$. Hence Q is not tied of type T1a. As $Z(G) = 1$, Q is not tied of type T1b. As u occurs as a letter in an element of S and $C_G(x) = C_G(S)$ it follows that $u \in A$, so Q is not tied of type T2. From Theorem 4.4, $\text{cdim}(G) = \text{cdim}(G_x)$.

□

Example 4.6. Using Theorem 4.4 we may compute the centraliser dimension of the semibraid groups. The n th *semibraid* group is defined to be the partially commutative group G_n with presentation

$$\langle x_1, \dots, x_n \mid [x_i, x_j] = 1, \text{ for all } i, j \text{ such that } |i - j| \geq 2 \rangle.$$

By inspection, using Proposition 3.3, it is found that the centraliser dimensions of the first few semibraid groups are $\text{cdim}(G_1) = 0$, $\text{cdim}(G_2) = \text{cdim}(G_3) = 2$ and $\text{cdim}(G_4) = \text{cdim}(G_5) = 4$. Proposition 3.3 also implies that $Z(G_n) = 1$, for all $n \geq 2$, as $C(x_1, \dots, x_n) = 1$. The following lemma shows that this pattern continues.

Lemma 4.7. *For $n \geq 0$, $\text{cdim}(G_{2n}) = \text{cdim}(G_{2n+1}) = 2n$.*

Proof. Let $n \geq 4$ and suppose that $\text{cdim}(G_n) = d$. Let Γ_{n+2} be the commutation graph of the group G_{n+2} . Let K be the partially group with commutation graph obtained from Γ_{n+2} by deleting the vertex x_{n+1} . Then $[x_j, x_{n+2}] = 1$, for $j = 1, \dots, n$, so it follows from Corollary 4.5.2b, with $S = 1$, that $\text{cdim}(K) = \text{cdim}(G_n)$. As $Z(G_{n+2}) = 1$ and $Z(K) = \langle x_{n+2} \rangle$ Corollary 4.5.1a implies that $\text{cdim}(G_{n+2}) = \text{cdim}(K) + 2 = d + 2$. Thus the lemma follows from the facts that $\text{cdim}(G_4) = \text{cdim}(G_5) = 4$. □

The example above shows that there are partially commutative groups of any given even centraliser dimension. It is well known that there are no groups of centraliser dimension 1 (see for example [7]). It was pointed out to us by K. Goda that there are no partially commutative groups of centraliser dimension 3 either. In fact if we suppose that such groups do exist then we can take G to be a partially commutative group of centraliser dimension 3 with the minimal number of vertices among all such groups. The centre of G will then be trivial (as otherwise we may remove all central generators). Now remove a vertex x from the commutation graph of G and let G_x be the corresponding group. Then $\text{cdim}(G_x) \neq 1$ and $\text{cdim}(G_x) \neq 3$, by minimality of G , so $\text{cdim}(G_x) = 2$. If $Z(G_x) \neq 1$ then Corollary 4.5.1a implies that $\text{cdim}(G) = 4$. Hence $Z(G_x) = 1$ and so, from [7, Proposition 3.9.1], G_x is a CT-group. It follows that G_x is a free product of free Abelian groups. Using Corollary 4.5 again it is not hard to see that the centraliser dimension of G must either be 2 or 4, a contradiction. Therefore there is no partially commutative group of centraliser dimension 3. However, as the following example shows, every odd number greater than 3 is the centraliser dimension of a partially commutative group.

Example 4.8. If G is a partially commutative group of centraliser dimension d then it follows from [14, Lemma 2.2.2] that the partially commutative group $G \times F$, where F is free of rank 2, has centraliser dimension $d + 2$. The group of Example 4.1 with commutation graph Γ_1 has centraliser dimension 5. Therefore there exist partially commutative groups of centraliser dimension $2m + 1$, for all $m \geq 2$.

References

- [1] F.A.M. Aldosray and I. Stewart, *Lie algebras with the minimal condition on centralizer ideals*, Hiroshima Math. J., **19** no. 2 (1989), 397–407
- [2] V. A. Antonov, *Groups with restriction on centralisers*, Chelyabinsk State Technical University, 1993, Part I: 147pp., Part II: 118pp., in Russian.
- [3] V. Bludov, *On locally nilpotent groups with the minimal condition on centralizers*. In *Groups St. Andrews 1997 in Bath, I*, London Math. Soc. Lecture Note Ser., **260**, 81–84, Cambridge Univ. Press, Cambridge, 1999.
- [4] R. M. Bryant and B. Hartley, *Periodic locally soluble groups with the minimal condition on centralisers*, J. Algebra, **61** (1979), 326–334.
- [5] R. M. Bryant, *Groups with the minimal condition on centralisers*, J. Algebra, **60** (1979), 371–383.
- [6] G. Duchamp and D. Krob, *Partially Commutative Magnus Transformations*, Internat. J. Algebra Comput. **3** No.1 (1993) 15–41.
- [7] A. J. Duncan, I. V. Kazachkov and V. N. Remeslennikov *Centraliser Dimension and Universal Classes of Groups*, submitted.

- [8] E.S. Esyp, I.V. Kazachkov and V.N. Remeslennikov, *Divisibility Theory and Complexity of Algorithms for Free Partially Commutative Groups*, In: Groups, Languages, Algorithms. Contemporary Mathematics **378** (2005) 319 – 348.
- [9] S. Humphries *On representations of Artin groups and the Tits conjecture* J. Algebra **169** no. 3 (1994) 847–862.
- [10] N. Ito, *On finite groups with given conjugate types. I* Nagoya Math. J. **6** (1953), 17–28.
- [11] O. H. Kegel and B. A. P. Wehrfritz, *Locally finite groups* Amsterdam-London, 1973.
- [12] L.F. Kosvincev, *Finite groups with maximal centralizers of their elements.* (Russian) Mat. Zametki **13** (1973), 577–580.
- [13] J. C. Lennox and J. E. Roseblade, *Centrality in Finitely Generated Soluble Groups*, J. Algebra **16** (1970),
- [14] A. Myasnikov and P. Shumyatsky, *Discriminating groups and c-dimension.* J. Group Theory, **7**, (2004), No. 1, 135–142.
- [15] R. Schmidt, *Zentralisatorverbände endlicher Gruppen*, Rend. Dem. Math. Univ. Padova, **44** (1970), 97–131.
- [16] R. Schmidt, *Characterisierung der einfachen Gruppe der Ordnung 175560 durch ihren Zentralisatorverbände*, Math. Z., **116** (1970), N 4, 299–306.
- [17] P. Vassiliou, *Eine Bemerkung über die von dem Zentralisator bestimmte Abbildung.* Prakt. Akad. Athēnōn **42** (1967), 286–289.
- [18] F.O. Wagner, *Nilpotency in groups with the minimal condition on centralizers*, J. Algebra **217** (1999), no. 2, 448–460.
- [19] B.A.F. Wehrfritz, *Remarks on centrality and cyclicity in linear groups*, J. Algebra **18** (1971) 229–236.
- [20] A. E. Zalesski, *Locally finite groups with the minimality condition for the centralisers*, Proceedings of Academy of Science of USSR, 1965, N3, 127–129, in Russian.

Andrew J. Duncan,
 School of Mathematics and Statistics,
 Merz Court, University of Newcastle,
 Newcastle upon Tyne, NE1 7RU, United Kingdom.
 e-mail: A.Duncan@ncl.ac.uk

Ilya V. Kazachkov
 Department of Mathematics and Statistics
 University of McGill

805 Sherbrooke West, Montreal,
Quebec, Canada, H3A 2K6.
e-mail: kazachkov@math.mcgill.ca

Vladimir N. Remeslennikov
e-mail: remesl@iitam.omsk.net.ru

Institute of Mathematics
(Siberian Branch of Russian Academy of Science)
Pevtsova st. 13
644099, Omsk, Russia,
+7 381 2 247041 Fax +7 381 2 234584

